

# CS TRACER BORON

10% W/V



## HOW IT WORKS

Tracer BORON is a novel amino acid chelated micronutrient concentrate of a highly soluble BORON solution which has been uniquely produced by blending amino acids, common metal chelating agents, complexing agents and an important micronutrient together in to a proprietary tiered chelation process. This stable chelated BORON product will address and positively reinforce BORON deficiency in growing plants to ensure physiological plant processes improve plant performance at harvest.

The improved bio-availability of Amino acid chelates can increase micronutrient (and macronutrient) uptake and availability markedly when compared to non chelated nutrient solutions commonly used in agricultural production systems. Also, when few trace element deficiencies have been detected in crops, any increases in yields may be due to a more regulated trace element balance and the symbiotic relationship between plants and amino acid derived chelates thus creating the ultimate synergy for optimum nutrient delivery and exchange.

## GETTING OPTIMUM RESULTS FROM TRACER BORON

For the best results Tracer BORON can be applied as a standard flowable solution at planting with most general fertilizer products or as a foliar applied product by boomspray during critical growth stages such as early emergence, mid tillering, stem elongation (depending on season and agronomy recommendations. Do not apply during or after flowering of any crop).

Tracer BORON applied on seed, in furrow or more importantly as a foliar spray, will provide an effective response and direct benefit to any plants indicating known and visible BORON trace element deficiency.

## DIRECTIONS

Agitate well before use. Store out of direct sunlight. It is expected that Tracer BORON will be compatible with most fertiliser products but any user always needs to conduct jar tests first to absolutely confirm.

## APPLICATION

### BROAD ACRE

Application Rate ► 0.2 L -2 L per/Ha

(Depending on advisor recommendation as BORON must only be used at low volumes per Hectare)

Best results will be obtained by 2 applications at the advised rate depending on crop growth stage. For extreme deficiencies or to respond to herbicide damage in growing crops then application of up to 3L per Ha may be required.

### ANALYSIS

CS TRACER BORON	W/V
Amino Nitrogen	1.76%
Magnesium	10%

(+/- 10% accuracy as per recognised guidelines)